



Report of consensus panel 1 from the 11th International Workshop on Waldenstrom's Macroglobulinemia on management of symptomatic, treatment-naïve patients

Christian Buske^{a,*}, Jorge J. Castillo^b, Jithma Prasad Abeykoon^c, Ranjana Advani^d, Suzanne O. Arulogun^e, Andrew R. Branagan^f, Xinxin Cao^g, Shirley D'Sa^e, Jian Hou^h, Prashant Kapoor^c, Efsthios Kastritisⁱ, Marie J. Kersten^j, Veronique LeBlond^k, Merav Leiba^l, Jeffrey V. Matous^m, Jonas Paludo^c, Lugui Qiuⁿ, Constantine S. Tam^o, Alessandra Tedeschi^p, Sheeba K. Thomas^q, Ibrahim Tohidi-Esfahani^r, Marzia Varettoni^s, Josephine M. Vos^j, Ramon Garcia-Sanz^t, Jesus San-Miguel^u, Meletios A. Dimopoulosⁱ, Steven P. Treon^b, Judith Trotman^r

^a University Hospital Ulm, Institute of Experimental Cancer Research, Ulm, Germany

^b Dana Farber Cancer Institute, Harvard Medical School, Boston MA

^c Mayo Clinic, Rochester MN

^d Stanford University Medical Center, Stanford CA

^e University College London Hospitals, London UK

^f Massachusetts General Hospital, Boston MA

^g Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

^h Renji Hospital, Shanghai China

ⁱ Department of Clinical Therapeutics, National and Kapodistrian University of Athens, Athens, Greece

^j Department of Hematology, Amsterdam UMC, University of Amsterdam, Cancer Center Amsterdam/LYMMCARE, Amsterdam, Netherlands

^k Groupe Hospitalier Pitié-Salpêtrière, Sorbonne University, Paris France

^l Faculty of Health Science, Ben-Gurion University of the Negev, Israel Assuta Ashdod University Hospital; Faculty of Health Science, Ben-Gurion University of the Negev, Israel Memorial Sloan Kettering Cancer Center, New York, NY

^m Colorado Blood Cancer Institute, Sarah Cannon Research Institute, Denver, CO

ⁿ National Clinical Medical Research Center for Blood Diseases, Institute of Hematology and Blood Diseases Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Tianjin, China

^o Alfred Health, Monash University, Melbourne, Australia

^p A.O. Ospedale Niguarda Ca' Granda, Milan, Italy

^q University of Texas, MD Anderson Cancer Center, Houston, TX

^r Concord Repatriation General Hospital, University of Sydney, Sydney, Australia

^s Division of Hematology, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy

^t Hematology Department, University Hospital of Salamanca, Research Biomedical Institute of Salamanca, CIBERONC and Center for Cancer Research-IBMCC (University of Salamanca-CSIC), Salamanca, Spain

^u Clínica Universidad de Navarra, Centro de Investigación Médica Aplicada, Instituto de Investigación Sanitaria de Navarra, Centro de Investigación Biomédica en Red Cáncer, Pamplona, Spain

ARTICLE INFO

Keywords:

Waldenstrom's Macroglobulinemia
Treatment-naïve
Chemoimmunotherapy
BTK-inhibitors
MYD88
CXCR4

ABSTRACT

Consensus Panel 1 (CP1) of the 11th International Workshop on Waldenstrom's Macroglobulinemia (IWWM-11) was tasked with updating guidelines for the management of symptomatic, treatment-naïve patients with WM. The panel reiterated that watchful waiting remains the gold standard for asymptomatic patients without critically elevated IgM or compromised hematopoietic function. For first-line treatment, chemoimmunotherapy (CIT) regimens such as dexamethasone, cyclophosphamide, rituximab (DRC), or bendamustine, rituximab (Benda-R) continue to play a central role in managing WM, as they are effective, of fixed duration, generally well-tolerated, and affordable. Covalent BTK inhibitors (cBTKi) offer a continuous, generally well-tolerated alternative for the primary treatment of

* Corresponding author: Christian Buske, MD, University Hospital Ulm, Institute of Experimental Cancer Research, Albert – Einstein Allee 11, 89081, Ulm, Germany.
E-mail address: christian.buske@uni-ulm.de (C. Buske).

WM patients, particularly those unsuitable for CIT. In a Phase III randomized trial updated at IWWM-11, the second-generation cBTKi, zanubrutinib, was less toxic than ibrutinib and induced deeper remissions, thus categorizing zanubrutinib as a suitable treatment option in WM. While the overall findings of a prospective, randomized trial updated at IWWM-11 did not show superiority of fixed duration rituximab maintenance over observation following attainment of a major response to Benda-R induction, a subset analysis showed benefit in patients >65 years and those with a high IPPSWM score. Whenever possible, the mutational status of *MYD88* and *CXCR4* should be determined before treatment initiation, as alterations in these 2 genes predict sensitivity towards cBTKi activity. Treatment approaches for WM-associated cryoglobulins, cold agglutinins, AL amyloidosis, Bing-Neel syndrome (BNS), peripheral neuropathy, and hyperviscosity syndrome follow the common principle of reducing tumor and abnormal protein burden rapidly and deeply to improve symptoms. In BNS, ibrutinib can be highly active and produce durable responses. In contrast, cBTKi are not recommended for treating AL amyloidosis. The panel emphasized that continuous improvement of treatment options for symptomatic, treatment-naïve WM patients critically depends on the participation of patients in clinical trials, whenever possible.

© 2023 Published by Elsevier Inc.

Introduction

In a modern era of widening global access to covalent Bruton tyrosine kinase inhibitors (cBTKi), several questions arise about using fixed-duration chemoimmunotherapy (CIT) regimens versus using these oral therapies of indefinite duration in the first-line treatment of WM. For those receiving rituximab-chemotherapy, we assess updated data on the 2 commonly used and highly effective regimens, Bendamustine-Rituximab (Benda-R) and Dexamethasone, Cyclophosphamide, and Rituximab (DRC), and review the role of rituximab maintenance. We discuss the patient populations in whom cBTKi might be suitable and preferred agents based on long-term data from phase II studies and a phase III randomized clinical trial comparing ibrutinib and zanubrutinib. We address the impact of our emerging understanding of genetic alterations of WM on treatment selection. We consider what agents to choose, and as importantly avoid, in specific first-line clinical situations. Even with these efficacious and well tolerated first-line treatment options, we reiterate the ongoing central role of watch-and-wait (W&W) for those with smoldering disease. As we move into a more “endemic” albeit still unpredictable phase of COVID-19, a nuanced, evidence-based approach to each patient with WM remains essential. Notwithstanding the data, cost implications result in differing access to WM therapies worldwide, and these first-line treatment guidelines endeavor to provide therapeutic options for all patients globally.

What are the preferable treatments for treatment naïve, symptomatic WM patients?

Rituximab-chemotherapy and the covalent BTK inhibitors (cBTKi), ibrutinib alone or with rituximab, and zanubrutinib are reasonable first-line treatment options for patients with WM [1]. A large, multicenter, retrospective study of treatment-naïve WM patients who received ibrutinib or Benda-R showed similar outcomes in PFS and overall survival with a median follow-up of 4.2 years suggesting that CIT with Benda-R or a cBTKi are similarly effective in treatment-naïve patients [2].

Chemoimmunotherapy

Benda-R is the CIT of choice for patients requiring rapid debulking or with a high tumor burden. No prospective data have compared Benda-R and DRC in WM. However, a retrospective analysis suggested a trend to longer PFS with Benda-R than DRC, and patient-derived data supports a longer time-to-next treatment (TTNT) with Benda-R [3,4]. For less fit patients with lower tumor burden, particularly those with a gradual progression to treatment

initiation, DRC is an effective and well-tolerated treatment option [5,6].

For patients treated with Benda-R, the pivotal STiL trial used six four-weekly cycles of bendamustine at a dose of 90 mg/m², demonstrating high efficacy and manageable toxicity for WM [7]. Doses as low as 70mg/m² and/or as few as 4 cycles may be sufficient in some cases [8]. However, no prospective study has analyzed 4 vs 6 cycles of Benda-R in WM.

Alternative regimens such as R-CHOP, R-CVP, fludarabine, or cladribine-based therapies are not preferred in WM because of higher toxicity, particularly neurological and hematological, without evidence of long-term efficacy benefit compared to Benda-R or DRC.

cBTKi

Where available, cBTKi should be preferentially used in patients with significant comorbidities or frailty predicting increased risk of chemotherapy-associated toxicity. In addition, cBTKi are an important treatment option in young patients, in whom alkylator exposure with its long-term consequences pertaining to second hematologic malignancies may not be desirable. However, there is no formal definition of what constitutes medically unfit patients in WM ineligible for CIT. Thus, the decision to offer a cBTKi instead of CIT must be made by each clinician based on individual patient characteristics. Individual co-morbidities should guide the use of first vs second-generation cBTKis (eg, cardiac toxicity of ibrutinib). The 1:1 randomized comparison between ibrutinib and zanubrutinib in patients with *MYD88*^{MUT} that was updated at IWWM-11 demonstrated a lower rate of atrial fibrillation (8% vs 25%) and hypertension (15% vs 26%) for zanubrutinib compared to ibrutinib [9,10]. A higher rate of neutropenia for zanubrutinib versus ibrutinib was also reported, but this did not translate into a higher rate of infections for patients on zanubrutinib with 45 months of median follow-up. Fewer patients in the zanubrutinib arm had an adverse event (AE) leading to treatment discontinuation (9% vs 20%) or dose reductions (14% vs 23%) [9,10]. This update charted a numerical trend of deeper, earlier, and more durable responses on zanubrutinib compared with ibrutinib [10]. Zanubrutinib provided faster and deeper responses in patients with *CXCR4*^{MUT}, and responses to zanubrutinib were also seen in a single arm observational cohort of *MYD88*^{WT} patients who showed continued deepening of categorical responses over time [10]. Overall, no difference in PFS or OS between the zanubrutinib and ibrutinib arms was observed, though patients with *CXCR4*^{MUT} disease showed a trend towards superior PFS with zanubrutinib [10]. Thus, both cBTKi are valid treatment options, and individual patient co-morbidities and *MYD88* and *CXCR4* mutation status can be considered when selecting between these 2 cBTKi. Phase II data on acalabrutinib suggest

Table 1

Selected data from prospective studies in treatment-naïve patients with Waldenström Macroglobulinemia.

Study	Regimen	N	PR or better	VGPR or better	PFS
Dimopoulos et al.[43]	Dexamethasone	72	74%	7%	35 mo (median)
Kastritis et al.[5]	Rituximab				
	Cyclophosphamide				
Rummel et al.[27,44]	Bendamustine	257	88%	4%	65 mo(median)
	Rituximab				
Treon et al.[45]	Bortezomib	23	83%	35%	66 mo(median)
Treon et al.[46]	Dexamethasone				
	Rituximab				
Dimopoulos et al.[47]	Bortezomib weekly	59	68%	10%	42 mo(median)
Gavriatopoulou et al.[18]	Dexamethasone				
	Rituximab				
Treon et al. [14]	Carfilzomib	28	68%	36%	46 mo (median)
Meid et al.[48]	Dexamethasone				
	Rituximab				
Castillo et al.[49]	Ixazomib	26	77%	19%	40 mo (median)
Castillo et al.[15]	Dexamethasone				
	Rituximab				
Buske et al.[7]	Bortezomib	102	81%	17%	81% at 24 mo
	Cyclophosphamide				
	Dexamethasone				
	Rituximab				
	Cyclophosphamide	100	70%	10%	73% at 24 mo
	Dexamethasone				
	Rituximab				
Treon et al.[50]	Ibrutinib	30	87%	30%	76% at 4 y
Castillo et al.[51]					
Dimopoulos et al.[52]	Ibrutinib	34	76%	27%	70% at 4.5 y
Buske et al.[53]	Rituximab				
	Rituximab	34	41%	9%	32% at 4.5 y
Tam et al.[9]	Zanubrutinib	19	74%	36%	78% at 42 mo
Dimopoulos et al.[10]					
	Ibrutinib	18	67%	22%	70% at 42 mo
Owen et al.[11]	Acalabrutinib	14	79%	NR	86% at 66 mo
Owen et al.[54]					

N = number of patients; NR = not reported; PFS = progression-free survival; PR = partial response; VGPR = very good partial response.

this cBTKi is also well tolerated in WM patients. However, only 14 patients were treated first-line in this trial with 106 patients [11] (Table 1).

Data on other second-generation cBTKis, orelabrutinib and tirabrutinib, in first-line treatment are limited, but show encouraging activity with sustained responses. Whenever possible, treatment with cBTKi should be continued until second-line treatment is initiated to minimise IgM rebound and “cytokine response” side effects that are seen in up to 20% of patients following withdrawal [12,13].

Is there a role for proteasome inhibitor-based therapy in treatment-naïve, symptomatic WM patients?

Bortezomib, the first in class proteasome inhibitor (PI), has shown activity in combination with rituximab with or without dexamethasone, or combined with DRC [6]. An early analysis of the ECWM Phase II comparison of B-DRC vs DRC in 202 patients showed that adding bortezomib was deliverable with a trend towards faster and deeper responses. However, there was increased neurotoxicity (18%) and a possible signal of increased infections. PFS, the primary study end point, was comparable between the quadruplet and DRC [7]. Carfilzomib or ixazomib in combination with rituximab and dexamethasone has shown high levels of activity and durability in treatment-naïve WM patients [14,15]. In front-line therapy, a PI-based regimen may be appropriate for patients with AL amyloidosis, light chain deposition disease, and renal compromise. While PIs may produce treatment-related neuropathy, subcutaneous bortezomib, carfilzomib or ixazomib may be less neuropathic in patients with peripheral neuropathy [16–18]. Carfilzomib should be avoided in patients with cardiopulmonary disease [14].

Is there a role for Rituximab monotherapy in treatment-naïve, symptomatic WM patients?

In 2023, there is a more limited role for rituximab monotherapy in the first-line treatment of WM, given major response rates of only 20% to 40% and median PFS of 16 to 18 months [19,20]. Patients being unfit for CIT may benefit from rituximab single agent treatment. Generally, however, if a patient is unfit for chemotherapy, a cBTKi is recommended. Single-agent rituximab may be an option in settings where cBTKi is unavailable. However, the time to response is long, the efficacy is moderate, and there is a risk of IgM flare in ~50% of patients [1,21,22]. Based on the latter, rituximab monotherapy should be avoided in patients with serum IgM levels >40 g/L, and apheresis facilities should be available for critically elevated IgM levels after treatment [22].

Role of COVID-19 and primary therapy of WM

As we move into the fourth year of the COVID-19 pandemic, it is recognized that WM patients show poor serological response to COVID-19 vaccination if they are on a cBTKi or had recent rituximab therapy [23–25]. Covid-specific T cell responses are preserved. Boosters may be of particular benefit in augmenting serological response to vaccination [25]. In this context, the impact of COVID-19 probably does not justify a compromise in therapy choice; patients should be offered the best treatment option, while optimizing immunization and pre-exposure prophylaxis in parallel, as well as education about the importance of seeking medical assessment and use of antiviral therapy, such as nirmatrelvir/ritonavir (Paxlovid). A discussion about the benefits and risks of treatment of COVID-19 infection should be considered for all patients, and in particular for those who are unvaccinated or who have not received updated

boosters for COVID-19 (see also CP5 report in this edition of Seminars in Hematology) [26].

Given the CYP3A inhibition of nirmatrelvir/ritonavir and the increase of the plasma concentration of cBTKis, a dose pause or 4-fold reduction in dosing is recommended while on and 48 hours immediately after the use of nirmatrelvir/ritonavir (see CP5 report in this edition of Seminars in Hematology). Both CIT and cBTKi are immunosuppressive and justify pre-exposure prophylaxis with anti-COVID-19 antibodies for WM patients, as long as variants remain sensitive to a given passive antibody combination.

In which patient should rituximab maintenance be used as part of frontline therapy of WM? Also, in light of the COVID-19 pandemic, should rituximab maintenance be held or shortened, and in what patient?

Based on prospective data, there is no general role for rituximab maintenance after first-line induction. In an update of the MAINTAIN study, which randomized >200 patients who attained at least a partial response to Benda-R induction to 2 years of maintenance vs observation, there was no significant difference in the PFS or OS between arms after a median of 7 years of follow-up. Of note, the median PFS in the observation arm was 84 months [27]. Based on these data, rituximab maintenance is not generally recommended for all WM patients who attained a major response after induction CIT. However, some WM patients may benefit since patients >65 years and those with high IPPSWM score showed an improved PFS, in a subset analysis. Maintenance rituximab after CIT may also be considered in patients with a minor response to induction or disease manifestations that might benefit from more profound responses (eg, renal involvement).

Which molecular characteristics have an impact on our treatment choice? MYD88? CXCR4? TP53?

The mutational status of MYD88, and increasingly CXCR4, guide treatment decisions in WM and should be determined whenever possible, particularly when considering BTK-inhibitor use (see above). Disruption or mutations of TP53 occur in up to 10% of treatment-naïve WM patients [28]. However, the precise impact of TP53 alterations on treatment outcomes in the primary treatment setting of WM is not currently known.

Additional questions submitted to consensus panel from the audience at IWWW-11

How should we manage asymptomatic treatment naïve patients in WM, including patients with high-risk disease?

W&W with serial evaluation of IgM levels, complete blood count, and patient symptoms remains the gold standard for asymptomatic patients with WM. Monitoring intervals should be individualized according to the dynamic of the disease in a given patient. The risk of progression can be estimated by a patient risk calculator ([Patient AWM Risk Calculator](#); available at www.awmrisk.com) dividing patients into high, intermediate, and low-risk groups with median estimated time to therapy of 1.8, 4.8, and 9.2 years, respectively [29]. Follow-up frequency can be adopted according to the risk of progression. Low-risk patients can be seen every 6 to 12 months, intermediate-risk every 4 to 6 months, and high-risk every 2 to 3 months. A high-risk category does not constitute a need to initiate treatment.

Should we use covalent BTK inhibitors for WM-related amyloidosis?

Data on the efficacy and safety of cBTKi in WM-related AL amyloidosis are scarce and limited to case series; these have demonstrated high toxicity rates and less efficacy [30,31]. In addition,

cBTKi should not be used in patients with cardiac amyloidosis, given the potential increased risk of arrhythmia. Thus, cBTKi are not the preferred treatment option for WM patients with AL amyloidosis (see CP6 report in this edition of Seminars in Hematology).

What is our treatment approach for cryoglobulins, cold agglutinins, amyloidosis, BNS, peripheral neuropathy and HV syndrome in WM?

Treatment approaches for WM-associated cryoglobulins, cold agglutinins, AL amyloidosis, Bing-Neel syndrome (BNS), peripheral neuropathy (PN), and hyperviscosity syndrome (HV) follow the common principle of reducing tumor burden and production of abnormal proteins as rapidly and deeply as possible. In BNS, ibrutinib can be highly active and produce durable responses [32]. In case of ibrutinib failure, CNS-penetrating compounds such as methotrexate, cytarabine, fludarabine and bendamustine show varying levels of activity [33]. Neurotoxic agents such as bortezomib, ixazomib, and vincristine should be avoided in WM-associated PN. Rituximab with or without chemotherapy could be an option according to the severity of anti-MAG neuropathy [34]. For symptomatic HV, plasmapheresis followed by systemic treatment is the recommended approach. WM patients with AL amyloidosis were shown to benefit from treatments including PIs or CIT such as Benda-R [35]. Cryoglobulins and cold agglutinins can be efficiently treated by CIT, PI-based treatment, or cBTKi [35].

Should the risk of therapy-associated secondary malignancies be factored in for selecting first-line treatment?

Several independent retrospective analyses have shown an increased risk for secondary malignancies (SM) in WM patients compared to the general population [36]. However, data quantifying the risk of therapy-induced SM in WM are still mounting. The risk of inducing SM by chemotherapy, especially treatment-related myeloid neoplasms and myelodysplasia, should be discussed with the patient. In addition, screening for SM should be part of the regular follow-up program in patients with WM (see also CP2 recommendations in this edition of Seminars in Hematology).

Should the “tempo of symptom progression” be factored in for decision-making on initiating treatment?

The dynamic of symptom progression is an important factor in tailoring follow-up intervals in the asymptomatic, W&W patients and the decision to initiate treatment. However, there is no formal assessment available categorizing the tempo of symptom progression. Thus, disease dynamics must be judged on a case-by-case basis. The decision to start, and choice of treatment should be based on individual patient characteristics, genomics, comorbidities, and disease dynamics.

When is plasmapheresis appropriate and how long should a patient be plasmapheresed?

Plasmapheresis is indicated in symptomatic hyperviscosity (HV) and should be considered in patients without symptomatic HV when a CD20-directed antibody containing regimen is planned for patients with serum IgM levels above 40 g/L to prevent an IgM flare [21,22]. Alternatively, to reduce the risk of IgM flare, the chemotherapy component may be administered without rituximab for 1 to 3 cycles until the IgM level is <40 g/L. Two to 3 plasmapheresis sessions, each performed ideally every other day can lower the IgM level by 30% to 60% and should continue until symptoms associated with HV are relieved. Plasmapheresis should always be followed by systemic “definitive” treatment as the effect of plasmapheresis on lowering IgM is transient. Avoidance of red

cell transfusions is recommended in this setting, and if necessary, this should be timed to follow plasmapheresis.

How should we “bridge” WM patients coming off BTK inhibitors?

An IgM rebound is a common occurrence following the discontinuation or withholding of a cBTKi [37]. Whenever possible, treatment with cBTKi should be continued until subsequent treatment is initiated and shows activity to avoid serum IgM rebound. Caution with even temporary combinations should be exercised, particularly with venetoclax, due to the unexpected risk of ventricular arrhythmias reported in combination with ibrutinib [38]. Oral prednisone might help mitigate the risk of withdrawal symptoms characterized by fever, body aches, night sweats, arthralgia, headache, and fatigue, which have been reported in approximately 20% of patients who stop cBTKi therapy [13]. The choice of subsequent therapy should be directed by the patient's symptoms, comorbidities, genomic profile, and previous therapy history (see also CP2 report in this edition of *Seminars in Hematology*). CIT, proteasome inhibitor-based therapy, venetoclax, noncovalent BTKi such as pirtobrutinib or clinical trials are reasonable options after progression on a cBTKi [12,39]. A switchover from ibrutinib or acalabrutinib to zanubrutinib without any bridge may also be considered for patients with intolerance to ibrutinib [40].

How should prognostic factors be incorporated into treatment recommendations?

The International Prognostic Scoring System for WM (IPSSWM) or revised IPSSWM (rIPSSWM) allows robust categorization of patients into different risk groups, separating patients with good prognosis from those with poor long-term outcomes. However, it is not recommended to stratify treatment according to the IPSSWM or rIPSSWM in routine clinical practice, as prospective data identifying the superiority of specific treatments in distinct risk groups is lacking. CXCR4 mutations and the lack of MYD88 mutations impact cBTKi efficacy and thus should be incorporated as predictive factors into treatment decisions (see above). TP53 mutations are associated with inferior outcomes, but prospective data analyzing their impact on outcome after CIT or cBTKi are lacking in the primary therapy of WM (see CP2 recommendations in this edition of *Seminars in Hematology* as well) [28].

Discussion

Since the publication of the last consensus treatment recommendations from the 10th IWWM, the dramatic impact of the COVID-19 pandemic reminds us of the vulnerability of WM patients caused by treatment-associated or disease-inherent immunosuppression. The quote of Jan Gosta Waldenström to “*let well alone*” remains central to our decisions to initiate treatment or to apply dose intense or continuous regimens. W&W remains the gold standard in 2023 in asymptomatic patients despite the emergence of well-tolerated chemotherapy-free approaches.

The class of cBTKi has gained relevance in the first-line treatment of WM, being now approved in many parts of the world as single agents or in combination with rituximab. We now have prospective, long-term data demonstrating that the second-generation cBTK-inhibitors zanubrutinib and acalabrutinib are highly effective in WM, with zanubrutinib showing less cardiac toxicity as compared to ibrutinib. However, the sequencing of cBTKi remains unclear, with scant data on the efficacy of subsequent CIT in patients who progress after first-line cBTKi.

CIT remains a central first-line treatment approach, and a common regimen used in the frontline setting in Europe, North America, Asia-Pacific and Latin America [2,41,42]. CIT has the advantage

over cBTKi of being a less costly fixed-duration treatment with the capacity to induce deep and durable remissions and comparable PFS to cBTKi in the case of Benda-R.[2] Prospective, randomized clinical trials comparing CIT with fixed-duration chemotherapy-free treatments are needed though to clarify the role of CIT in an era with rapidly emerging novel treatment concepts.

Although there is consensus that the mutational status of MYD88 and CXCR4 should be determined in all WM patients, technical challenges in identifying CXCR4 mutations and the relative impact of different CXCR4 mutations hinder the introduction of widespread, reliable testing in clinical practice and a genotype-driven therapeutic approach in WM. It is also accepted that other molecular alterations in WM, such as TP53, will likely impact treatment outcome, but little prospective data currently exists on the impact of these on treatment response and duration in the primary therapy of WM.

We appreciate the extraordinary progress made in the initial clinical management of WM since our last consensus statement. Expanded first-line treatment options are ensuring more patients achieve longer first remissions. Nonetheless, major challenges lie ahead for our ambitious goal to establish well-tolerated, molecularly and clinically guided personalised fixed-duration chemotherapy-free approaches to ensure long-lasting quality remissions. The greatest challenge will be to ensure such approaches are accessible and affordable for all patients globally. Finally, we advocate for the thoughtful design of clinical trials in the frontline setting and strongly encourage patient participation.

Authors' Contributions

CB, JT, JJC, RGS, JSM, and SPT prepared, reviewed and submitted key questions for Consensus Panel 1 (CP1). Questions were reviewed in an open general assembly by attendants of IWWM-11, and additional questions for CP1 deliberations were formulated and submitted. CB wrote the first draft of CP1 responses, and draft was reviewed and modified by CB, JT, JJC, RGS, JSM, and SPT. Final draft was submitted to CP1 general panel for review and commentary. CP1 general panel was composed of individuals with experience in the care of WM patients who attended IWWM-11 and volunteered to be on CP1 panel.

Disclosure

CB received honoraria from Roche, Pfizer, Janssen, Hexal, Celltrion, AbbVie, Novartis, Bayer, Morphosys, Regeneron, Beigene; consulting fees from Roche, Pfizer, Janssen, Hexal, Celltrion, AbbVie, Novartis, Bayer, Morphosys, Regeneron, Beigene, Sobi; and Research Funding from Roche, Janssen, Celltrion, AbbVie, Bayer, Amgen, and MSD.

JJC received research funds from Abbvie, AstraZeneca, Beigene, Cellectar, LOXO, Pharmacyclics, TG Therapeutics, and honoraria from Abbvie, Beigene, Cellectar, Kite, LOXO, Janssen, Pharmacyclics, and Roche Pharmaceuticals.

SPT received research funding, and/or consulting fees from Abbvie/Pharmacyclics Inc., Janssen Oncology Inc., Beigene Inc., Eli Lilly Pharmaceuticals, and Bristol Myers Squibb.

JSM declares participation on advisory boards and consulting services, on behalf of my Institution, for Abbvie, Amgen, BMS, Celgene, GSK, Haemalogix, Janssen-Cilag, Karyopharm, MSD, Novartis, Pfizer, Takeda, Regeneron, Roche, Sanofi, and SecuraBio.

JT receives research funds to her institution from Beigene, Pharmacyclics, Janssen, Roche, Cellectar, Takeda and BMS.

LQ declares participation on advisory boards and consulting services for Beigene, Xi'an Janssen, Pfizer, Sanofi, AstraZeneca

JP declares honoraria from AbbVie and research funding to his institution from Biofourmis and Karyopharm.

VL received consulting and advisory boards participation fees from Abbvie, MSD, Janssen-Cilag, Astra-Zeneca, Amgen Beigene Lilly.

ARB reports being and advisory board member for Adaptive, Beigene, CSL Behring, Genzyme, Karyopharm, Pharmacyclics, and Sanofi.

PK received research funding from Amgen, Regeneron, Bristol Myers Squibb, Loxo Pharmaceuticals, Ichnos, Karyopharm, Sanofi, AbbVie and GlaxoSmithKline; Honoraria from BeiGene, Pharmacyclics, X4 Pharmaceuticals, Oncopptides, Angitia Bio, Glaxo-SmithKline, AbbVie and Sanofi.

SD received research funding, and/or consulting fees from BeiGene and Janssen and Kite Pharma.

EK declares honoraria from Amgen, Janssen, GSK, Pfizer and research funding from Amgen, Janssen, GSK, and Pfizer.

JVM has served on advisory committee's for Pharmacyclics, Janssen and BeiGene.

RGS declares honoraria from Amgen, Takeda, Janssen, Incyte, Astellas, BeiGene, AstraZeneca, Pfizer; and research funding from Novartis, Gilead, Astellas, Janssen; and participation in advisory boards for Amgen, Pharmacyclics, Takeda.

CST reports research funding from Janssen, AbbVie and Beigene; and honoraria from Janssen, Pharmacyclics, Beigene and AbbVie.

MJK reports honoraria from Kite, Novartis, and Miltenyi Biotech, Roche, and Bristol Myers Squibb/Celgene; consultancy or advisory role for Kite, Roche, Bristol Myers Squibb/Celgene, Novartis, Adicet Bio and Miltenyi Biotech; research funding from Kite; and travel support from Kite, Roche, Novartis, and Miltenyi Biotech.

SKT reports research funding from Collectar Biosciences, X4 Pharma, Ascentage Pharma, Bristol Myers Squibb, Acerta Pharma, Genentech and Consulting for Collectar Biosciences.

RA reports institute research funding: Seattle Genetics, Pharmacyclics, Regeneron, Gilead, Merck, Cyteir, ADCT, Daiichi Sanyo. Advisory Board: Roche, BMS, Beigene, ADCT.

MV received honoraria from Abbvie, AstraZeneca, Beigene and Janssen-Cilag.

AT declares being on the Advisory Board and Speaker Bureau for Janssen SPA, AbbVie, Astrazeneca, and Beigene.

JMIV received institutional honoraria for research support (Beigene, Abbvie/Genmab); Advisory Board/consultancy (Sanofi); Speakers buro (BMS, Sanofi).

MAD received honoraria from Amgen, Bristol Myers Squibb, GSK, Janssen, Beigene Inc, Sanofi and Takeda.

SOA has no declarations.

ITE has no declarations.

JPA has no disclosures.

ZC has no disclosures.

JH has no disclosures.

ML has no disclosures.

Acknowledgments

The authors gratefully acknowledge Beigene Pharmaceuticals, Abbvie/Pharmacyclics, Janssen Pharmaceuticals, the International Waldenström's Macroglobulinemia Foundation, and Collectar Biosciences, Inc. for their support of the 11th International Workshop on Waldenström's Macroglobulinemia. Consensus panel reports of IWWM-11 are for educational purposes and should not be construed as offering specific medical advice for patients.

References

- [1] Dimopoulos MA, kastritis E, Owen RG, et al. Treatment recommendations for patients with Waldenström Macroglobulinemia and related disorders: IWWM-7 consensus. *Blood* 2014;124:1404–11.
- [2] Abeykoon JP, Kumar S, Castillo JJ, et al. Bendamustine rituximab versus ibrutinib as primary therapy for Waldenström Macroglobulinemia: an International Collaborative Study. *J Clin Oncol* 2022;40(16):7566.
- [3] Abeykoon JP, Zanwar S, Ansell SM, et al. Assessment of fixed-duration therapies for treatment-naïve Waldenström Macroglobulinemia. *Ann Hematol* 2021;96(8):945–53.
- [4] Tohidi-Esfahani I, Warden A, Malunis E, et al. Whimsical: a global Waldenström's Macroglobulinemia patient-derived data registry capturing treatment and quality of life outcomes. *Am J Hematol* 2021;96:e218–22.
- [5] Rummel MJ, Niederle N, Maschmeyer G, et al. Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet* 2013;381:1203–10.
- [6] Kastritis E, Gavriatopoulou M, Kyrtsonis MC, et al. Dexamethasone, rituximab, and cyclophosphamide as primary treatment of Waldenström Macroglobulinemia: final analysis of a phase 2 study. *Blood* 2015;126:1392–4.
- [7] Buske C, Dimopoulos MA, Grunenberg A, et al. Bortezomib-dexamethasone, rituximab, and cyclophosphamide as first-line treatment for Waldenström's Macroglobulinemia: a prospectively randomized trial of the European Consortium for Waldenström's Macroglobulinemia. *J Clin Oncol* 2023;JCO2201805 (Still ahead of print).
- [8] Cheson BD, Brugger W, Damaj G, et al. Optimal use of bendamustine in hematologic disorders: treatment recommendations from an international consensus panel—an update. *Leuk lymphoma* 2016;57:766–82.
- [9] Tam CS, Opat S, D'sa S, et al. A randomized phase 3 trial of zanubrutinib vs ibrutinib in symptomatic Waldenström Macroglobulinemia: the ASPEN study. *Blood* 2020;136:2038–50.
- [10] Dimopoulos M, Opat S, D'sa S, et al. ASPEN: long-term follow-up results of a phase 3 randomized trial of zanubrutinib vs ibrutinib in patients with waldenström macroglobulinemia (wm). In: *Proc. Of the 11th International Workshop on Waldenström Macroglobulinemia*; 2022. p. 39.
- [11] Owen RG, McCarthy H, Rule S, et al. Acalabrutinib monotherapy in patients with Waldenström Macroglobulinemia: a single-arm, multicentre, phase 2 study. *Lancet Haematol* 2020;7:e112–21.
- [12] Gustine JN, Sarosiek S, Flynn CA, et al. Natural history of Waldenström Macroglobulinemia following acquired resistance to ibrutinib monotherapy. *Haematologica* 2022;107:1163–71.
- [13] Castillo JJ, Gustine JN, Meid K, et al. Ibrutinib withdrawal symptoms in patients with Waldenström Macroglobulinemia. *Haematologica* 2018;103:e307–10.
- [14] Treon SP, Tripsas CK, Meid K, et al. Carfilzomib, rituximab, and dexamethasone (CARD) treatment offers a neuropathy-sparing approach for treating Waldenström's Macroglobulinemia. *Blood* 2014;124:503–10.
- [15] Castillo JJ, Meid K, Flynn CA, et al. Ixazomib, dexamethasone, and rituximab in treatment-naïve patients with Waldenström Macroglobulinemia: long-term follow-up. *Blood Adv* 2020;4:3952–9.
- [16] Moreau P, Pylypenko H, Grosicki S, et al. Subcutaneous versus intravenous administration of bortezomib in patients with relapsed multiple myeloma: a randomised, phase 3, non-inferiority study. *Lancet Oncol* 2011;12(5):431–40.
- [17] Ghobrial IM, Xie W, Padmanabhan S, et al. Phase II trial of weekly bortezomib in combination with rituximab in untreated patients with Waldenström Macroglobulinemia. *Am J Hematol* 2010;85:670–4.
- [18] Gavriatopoulou M, García-Sanz R, Kastritis E, et al. BDR in newly diagnosed patients with WM: Final analysis of a phase 2 study after a minimum follow-up of 6 years. *Blood* 2017;129:456–9.
- [19] Dimopoulos MA, Zervas C, Zomas A, et al. Treatment of Waldenström's Macroglobulinemia with rituximab. *J Clin Oncol* 2002;20:2327–33.
- [20] Treon SP, Emmanouilides C, Kimby E, et al. Extended rituximab therapy in Waldenström's Macroglobulinemia. *Ann Oncol* 2005;16(1):132–8.
- [21] Ghobrial IM, Fonseca R, Greipp PR, et al. Initial Immunoglobulin M 'flare' after rituximab therapy in patients diagnosed with Waldenström Macroglobulinemia: an Eastern Cooperative Oncology Group study. *Cancer* 2004;101:2593–8.
- [22] Treon SP, Branagan AR, Hunter ZR, et al. Paradoxical increases in serum IgM and viscosity levels following rituximab in Waldenström's Macroglobulinemia. *Ann Oncol* 2004;15(10):1481–3.
- [23] Gavriatopoulou M, Terpos E, Ntanasis-Stathopoulos I, et al. Poor neutralizing antibody responses in 106 patients with WM after vaccination against Sars-Cov-2: a prospective study. *Blood Adv* 2021;5:4398–405.
- [24] Rosati M, Terpos E, Bear J, et al. Low spike antibody levels and impaired ba.4/5 neutralization in patients with multiple myeloma or Waldenström's Macroglobulinemia after BNT162b2 booster vaccination. *Cancers (basel)* 2022;14:5816.
- [25] Branagan AR, Lei MM, Mo CC, et al. Clinical effectiveness and long-term serologic responses of COVID-19 vaccination in patients with multiple myeloma and Waldenström Macroglobulinemia. *Blood* 2022;140:10094–6.
- [26] Talaullikar D, Advani RH, Branagan AR, et al. Consensus statement on the management of Waldenström Macroglobulinemia patients during the COVID-19 pandemic. *Hemasphere* 2020;4:e433.
- [27] Rummel M. Presentation of STIL Maintenance Study of Rituximab in Waldenström Macroglobulinemia. In: *Proc. 11th International Workshop on WM*, Madrid, Spain; 2022.
- [28] Gustine JN, Tsakmaklis N, Demos MG, et al. TP53 mutations are associated with mutated MYD88 and CXCR4, and confer an adverse outcome in Waldenström Macroglobulinemia. *Br J Haematol* 2019;184:242–5.
- [29] Bustoros M, Sklaventis-Pistofidis R, Kapoor P, et al. Progression risk stratification of asymptomatic Waldenström Macroglobulinemia. *J Clin Oncol* 2019;37:1403–11.
- [30] Pika T, Hegenbart U, Flodrova P, et al. First report of ibrutinib in IgM-related amyloidosis: few responses, poor tolerability, and short survival. *Blood* 2018;131:368–71.

- [31] Gustine JN, Szalat RE, Staron A, et al. Light chain amyloidosis associated with Waldenstrom Macroglobulinemia: treatment and survival outcomes. *Haematologica* 2022 (Still ahead of print). doi:10.3324/haematol.2022.282264.
- [32] Castillo JJ, Itchaki G, Paludo J, et al. Ibrutinib for the treatment of Bing-Neel Syndrome: a multicenter study. *Blood* 2019;133:299–305.
- [33] Castillo JJ, D'Sa Lunn MP, et al. Central nervous system involvement by Waldenstrom macroglobulinemia (Bing Neel Syndrome): a multiinstitutional retrospective study. *Br J Haem* 2016;172(5):709–15.
- [34] Nivet T, Baptiste A, belin L, et al. Immunochemotherapy versus rituximab in anti-myelin-associated glycoprotein neuropathy: a report of 64 patients. *Br J Haematol* 2022;198(2):298–306.
- [35] Dimopoulos MA, Kastritis E. How I treat Waldenstrom Macroglobulinemia. *Blood* 2019;134:2022–35.
- [36] Castillo JJ, Olszewski AJ, Hunter ZR, et al. Incidence of secondary malignancies among patients with Waldenstrom Macroglobulinemia: an analysis of the SEER database. *Cancer* 2015;121:2230–6.
- [37] Gustine JN, Meid K, Dubeau T, et al. Ibrutinib discontinuation in Waldenstrom Macroglobulinemia: etiologies, outcomes, and IgM rebound. *Am J Hematol* 2018;93:511–17.
- [38] Castillo JJ, Sarosiek SR, Branagan AR, et al. Ibrutinib and venetoclax in previously untreated Waldenstrom Macroglobulinemia. *Proc Am Soc Hematology* 2022;231.
- [39] Palomba ML, Patel MR, Eyre TA, et al. Efficacy of pirtobrutinib, a highly selective, non-covalent (reversible) BTK inhibitor in relapsed /refractory waldenström macroglobulinemia: results from the phase 1/2 BRUIN study. *Proc Am Soc Hematology* 2022;229.
- [40] Shadman M, Flinn IW, Moshe Y, et al. Zanubrutinib in patients with previously treated B-cell malignancies intolerant of previous Bruton tyrosine kinase inhibitors in the USA: a phase 2, open-label, single-arm study. *Lancet Haematol* 2023;10(1):e35–45.
- [41] Buske C, Sadullah S, Kastritis E, et al. Treatment and outcome patterns in European patients with Waldenstrom's Macroglobulinaemia: a large, observational, retrospective chart review. *Lancet Haematol* 2018;5:e299–309.
- [42] Riva E, Duarte PJ, Valcarcel B, et al. Treatment and survival outcomes of Waldenstrom Macroglobulinemia in Latin American patients: a multinational retrospective cohort study. *JCO Glob Oncol* 2022;8:e2100380.
- [43] Dimopoulos MA, Anagnostopoulos A, Kyrtonis MC, et al. Primary treatment of Waldenstrom Macroglobulinemia with dexamethasone, rituximab, and cyclophosphamide. *J Clin Oncol* 2007;25:3344–9.
- [44] Rummel MJ, Lerchenmüller C, Hensel M, et al. Two years rituximab maintenance vs. observation after first line treatment with bendamustine plus rituximab in patients with Waldenström's Macroglobulinemia: results of a prospective, randomized, multicenter phase 3 study (the STIL NHL 7-2008 MAINTAIN Trial). *Blood* 2019;134:343.
- [45] Treon SO, Ioakimidis, Soumerai JD, et al. Primary therapy of Waldenstrom Macroglobulinemia with bortezomib, dexamethasone, and rituximab: WMCTG clinical trial 05-180. *J Clin Oncol* 2009;27:3830–5.
- [46] Treon SP, Meid K, Gustine J, et al. Long-term outcome of a prospective study of bortezomib, dexamethasone and rituximab (BDR) in previously untreated, symptomatic patients with Waldenström's Macroglobulinemia. *Blood* 2015;126:1833.
- [47] Dimopoulos MA, Garcia-Sanz R, Gavriatopoulou M, et al. Primary therapy of Waldenstrom Macroglobulinemia with weekly bortezomib, low-dose dexamethasone, and rituximab (BDR): long-term results of a phase 2 study of the European Myeloma Network (EMN). *Blood* 2013;122:3276–82.
- [48] Meid K, Dubeau T, Severns P, et al. Long-term follow-up of a prospective clinical trial of carfilzomib, rituximab and dexamethasone (CARD) in Waldenstrom's macroglobulinemia. *Blood* 2017;130:2772.
- [49] Castillo JJ, Meid K, Gustine JN, et al. Prospective clinical trial of ixazomib, dexamethasone, and rituximab as primary therapy in Waldenstrom Macroglobulinemia. *Clin Cancer Res* 2018;24:3247–52.
- [50] Treon SP, Gustine J, Meid K, et al. Ibrutinib monotherapy in symptomatic, treatment-naïve patients with Waldenstrom Macroglobulinemia. *J Clin Oncol* 2018;36:2755–61.
- [51] Castillo JJ, Meid K, Gustine JN, et al. Long-term follow-up of ibrutinib monotherapy in treatment-naïve patients with waldenstrom macroglobulinemia. *Leukemia* 2022;36:532–9.
- [52] Dimopoulos MA, Tedeschi A, Trotman J, et al. Phase 3 trial of ibrutinib plus rituximab in waldenstrom's macroglobulinemia. *N Engl J Med* 2018;378:2399–410.
- [53] Buske C, Tedeschi A, Trotman J, et al. Ibrutinib plus rituximab versus placebo plus rituximab for Waldenstrom's Macroglobulinemia: final analysis from the randomized phase III INNOVATE study. *J Clin Oncol* 2022;40:52–62.
- [54] Owen R, McCarthy H, Rule S, et al. Acalabrutinib in treatment-naïve or relapsed/refractory Waldenström Macroglobulinemia: 5-year follow-up of a phase 2, single-arm study. *HemaSphere* 2022;6:1020–1.